

## PFKFB1/4 Blocking Peptide

Catalog #	Source	Reactivity	Applications
CBP1875	Synthetic	H, M, R, B, C, Mk, Z	BL
<b>Description</b>	The peptide is used to block Anti-PFKFB1/4 Antibody (#CPA1875) reactivity.		
<b>Form</b>	Lyophilized powder		
<b>Gene Symbol</b>	PFKFB1; PFKFB4		
<b>Alternative Names</b>	PFKFB1; F6PK; PFRX; 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 1; 6PF-2-K/Fru-2,6-P2ase 1; PFK/FBPase 1; 6PF-2-K/Fru-2,6-P2ase liver isozyme; PFKFB4; 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 4; 6PF-2-K/Fru-2,6-P2ase 4; PFK/FBPase 4; 6PF-2-K/Fru-2,6-P2ase testis-type isozyme		
<b>Entrez Gene</b>	5207, 5210 (Human); 18639, 270198 (Mouse); 24638, 54283 (Rat)		
<b>SwissProt</b>	P16118, Q16877 (Human); P70266, Q6DTY7 (Mouse); P07953, P25114 (Rat)		
<b>Purity</b>	>85%		
<b>Quality Control</b>	The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry.		
<b>Directions for Use</b>	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours.		
<b>Storage/Stability</b>	Shipped at 4°C. Store at -20°C for one year.		

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

**WEB**  
www.cohesionbio.com

**ORDER**  
order@cohesionbio.com

**SUPPORT**  
techsupport@cohesionbio.com

**CUSTOM**  
custom@cohesionbio.com