

Product Data Sheet

PKC gamma Blocking Peptide

| Catalog # | Source | Reactivity | Applications | | | |
|----------------|------------|---|---|--|--|--|
| CBP1936 | Synthetic | H, M, R, B, Mk, Rb | BL | | | |
| Description | The | peptide is used to block A | Anti-PKC gamma Antibody (#CPA1936) reactivity. | | | |
| Form | Lyop | Lyophilized powder | | | | |
| Gene Symbol | PRKC | PRKCG | | | | |
| Alternative N | ames PKCC | PKCG; Protein kinase C gamma type; PKC-gamma | | | | |
| Entrez Gene | 5582 | 5582 (Human); 18752 (Mouse); 24681 (Rat) | | | | |
| SwissProt | P051 | P05129 (Human); P63318 (Mouse); P63319 (Rat) | | | | |
| Purity | >85% | >85% | | | | |
| Quality Contr | ol The | The quality of the peptide was evaluated by reversed-phase HPLC and by mass | | | | |
| | spec | trometry. | | | | |
| Directions for | Use Block | Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to | | | | |
| | antik | oody) and incubate the m | nixture at 4°C for overnight or at room temperature for | | | |
| | 2 ho | urs. | | | | |
| Storage/Stab | ility Ship | oed at 4°C. Store at -20°C | C for one year. | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |