

## **Product Data Sheet**

## PKA C gamma Blocking Peptide

Catalog # Source Reactivity Applications

CBP4168 Synthetic H BL

**Description** The peptide is used to block Anti-PKA C gamma Antibody (#CPA4168) reactivity.

Form Lyophilized powder

Gene Symbol PRKACG

Alternative Names cAMP-dependent protein kinase catalytic subunit gamma; PKA C-gamma

Entrez Gene 5568 (Human)

SwissProt P22612 (Human)

Purity >85%

Quality Control The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

**Directions for Use** Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

**Storage/Stability** Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com