

Product Data Sheet

Histone Deacetylase 2 (pS394) Blocking Peptide

Catalog #	Source	Reactivity	Applications
CBP5764	Synthetic	H, M, R, C	BL
Description	The peptide is used to block Anti-Histone Deacetylase 2 (pS394) Antibody (#CPA5764) reactivity.		
Form	Lyophilized powder		
Gene Symbol	HDAC2		
Alternative Names	Histone deacetylase 2; HD2		
Entrez Gene	3066 (Human); 15182 (Mouse)		
SwissProt	Q92769 (Human); P70288 (Mouse)		
Purity	>85%		
Quality Control	The quality of the peptide was evaluated by reversed-phase HPLC and by mass spectrometry.		
Directions for Use	Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to antibody) and incubate the mixture at 4°C for overnight or at room temperature for 2 hours.		
Storage/Stability	Shipped at 4°C. Store at -20°C for one year.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com