

## **Product Data Sheet**

## **HMGN2 (AcK31) Blocking Peptide**

Catalog # Source Reactivity Applications

CBP5849 Synthetic H, M, R, B, C, D, P BI

Description The peptide is used to block Anti-HMGN2 (AcK31) Antibody (#CPA5849) reactivity.

Form Lyophilized powder

Gene Symbol HMGN2

Alternative Names HMG17; Non-histone chromosomal protein HMG-17; High mobility group

nucleosome-binding domain-containing protein 2

Entrez Gene 3151 (Human); 15331 (Mouse)

SwissProt P05204 (Human); P09602 (Mouse); P18437 (Rat)

Purity >85%

Quality Control The quality of the peptide was evaluated by reversed-phase HPLC and by mass

spectrometry.

**Directions for Use** Blocking Peptide to the diluted primary antibody in a molar ratio of 10:1 (peptide to

antibody) and incubate the mixture at 4°C for overnight or at room temperature for

2 hours.

**Storage/Stability** Shipped at 4°C. Store at -20°C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com