

Product Data Sheet

Anti-CD22 Antibody-APC labled

| Catalog # | Source | e Reactivity | Applications | | | |
|--------------------|--------|----------------------------------------------------------------------------------------|-------------------------------------------------------|--|--|--|
| CFB8986 | Rat | М | IF, FC | | | |
| Description | | Rat monoclonal antibody APC labled to CD22 | | | | |
| Immunogen | | Splenic lymphocyte plasma | membranes from CBA x C57 F1 mice | | | |
| Purification | | The antibody was purified b | y affinity chromatography. | | | |
| Specificity | | Recognizes mouse CD22 | | | | |
| Clonality | | Monoclonal (clone: 2D6) | | | | |
| Conjugation | | APC | | | | |
| Form | | Rat IgG1 kappa. Liquid in PBS, pH 7.3, and 0.02% sodium azide. | | | | |
| Dilution | | 10 μl / assay | | | | |
| Gene Symbol | | CD22 | | | | |
| Alternative Names | | SIGLEC2; B-cell receptor CD22; B-lymphocyte cell adhesion molecule; BL-CAM; Sialic | | | | |
| | | acid-binding Ig-like lectin 2; | Siglec-2; T-cell surface antigen Leu-14; CD22 | | | |
| Entrez Gene | | 12483 (Mouse) | | | | |
| SwissProt | | P35329 (Mouse) | | | | |
| Directions for Use | | 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5 | | | | |
| | | ml tube. | | | | |
| | | 2. Add 10 μl labeled antiboo | ly to the bottom of flow tube mixing with the whole | | | |
| | | blood, incubate for 20 minu | tes at room temperature away from light. | | | |
| | | 3. Add 2 ml RBC lysis buffer, | incubate for 10 minutes away from light after mixing, | | | |
| | | dissolve red blood cells. | | | | |
| | | 4. Sample tube is set to 100 | 0 rpm centrifugation for 5 minutes, discard the | | | |
| | | supernatant. | | | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com