

# **Product Data Sheet**

# Anti-CD19 Antibody-APC/Cy5.5 labled

Catalog #	Source	Reactivity	Applications		
CFC8780	Mouse	M	IF, FC		
Description		Mouse monoclonal antiboo	ly APC/Cy5.5 labled to CD19		
Immunogen		Mouse CD19-expressing K5	62 human erythroleukemia cells		
Purification		The antibody was purified I	by affinity chromatography.		
Specificity		Recognizes mouse CD19			
Clonality		Monoclonal (clone: MB19-:	1)		
Conjugation		APC/Cy5.5			
Form		Mouse IgA kappa. Liquid in	PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution		10 µl / assay			
Gene Symbol		CD19			
Alternative Names		B-lymphocyte antigen CD19	9; B-lymphocyte surface antigen B4; Differentiation		
		antigen CD19; T-cell surface	e antigen Leu-12; CD19		
Entrez Gene		12478 (Mouse)			
SwissProt		P25918 (Mouse)			
Directions for Use		1. Take 100 μl peripheral bl	ood anticoagulated by EDTA and add to the bottom of 5		
		ml tube.			
		2. Add 10 $\mu l$ labeled antibo	dy to the bottom of flow tube mixing with the whole		
		blood, incubate for 20 minu	ites at room temperature away from light.		
		3. Add 2 ml RBC lysis buffer	, incubate for 10 minutes away from light after mixing,		
		dissolve red blood cells.			
		4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the		
		supernatant.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com