

Product Data Sheet

Anti-CD117 Antibody-APC/Cy7 labled

CFD8487MouseHIF, FCDescriptionMouse monoclonal antibody APC/Cy7 labled to CD117ImmunogenNative purified human CD117.PurificationThe antibody was purified by affinity chromatography.SpecificityRecognizes human CD117ClonalityMonoclonal (clone: 104D2)ConjugationAPC/Cy7FormMouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.Dilution10 µl / assayGene SymbolKITAlternative NameSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Catalog #	Source	e Reactivity	Applications			
ImmunogenNative purified human CD117.PurificationThe antibody was purified by affinity chromatography.SpecificityRecognizes human CD117ClonalityMonoclonal (clone: 104D2)ConjugationAPC/Cy7FormMouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.Dilution10 µl / assayGene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5	CFD8487	Mouse	e H	IF, FC			
PurificationThe antibody was purified by affinity chromatography.SpecificityRecognizes human CD117ClonalityMonoclonal (clone: 104D2)ConjugationAPC/Cy7FormMouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.Dilution10 µl / assayGene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Description	·					
SpecificityRecognizes human CD117ClonalityMonoclonal (clone: 104D2)ConjugationAPC/Cy7FormMouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.Dilution10 μl / assayGene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for UseI. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Immunogen		Native purified human CD117.				
ClonalityMonoclonal (clone: 104D2)ConjugationAPC/Cy7FormMouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.Dilution10 μl / assayGene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for UseI. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Purification		The antibody was purified by affinity chromatography.				
ConjugationAPC/Cy7FormMouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.Dilution10 μl / assayGene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for UseI. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Specificity		Recognizes human C	D117			
FormMouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.Dilution10 μl / assayGene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for UseI. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Clonality		Monoclonal (clone: 1	L04D2)			
Dilution10 μl / assayGene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Conjugation		APC/Cy7				
Gene SymbolKITAlternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Form		Mouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Alternative NamesSCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT; Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Dilution		10 μl / assay				
 Proto-oncogene c-Kit; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman 4 feline sarcoma viral oncogene homolog; CD antigen CD117 Entrez Gene 3815 (Human) SwissProt P10721 (Human) Directions for Use 1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 	Gene Symbol		КІТ				
4 feline sarcoma viral oncogene homolog; CD antigen CD117Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Alternative Names		SCFR; Mast/stem cell growth factor receptor Kit; SCFR; Piebald trait protein; PBT;				
Entrez Gene3815 (Human)SwissProtP10721 (Human)Directions for Use1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5			Proto-oncogene c-Kit	t; Tyrosine-protein kinase Kit; p145 c-kit; v-kit Hardy-Zuckerman			
SwissProtP10721 (Human)Directions for Use1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5			4 feline sarcoma vira	l oncogene homolog; CD antigen CD117			
Directions for Use 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5	Entrez Gene		3815 (Human)				
	SwissProt		P10721 (Human)				
	Directions for Use		1. Take 100 μl periph	eral blood anticoagulated by EDTA and add to the bottom of 5			
ml tube.			ml tube.				
2. Add 10 μl labeled antibody to the bottom of flow tube mixing with the whole			2. Add 10 µl labeled	antibody to the bottom of flow tube mixing with the whole			
blood, incubate for 20 minutes at room temperature away from light.			blood, incubate for 2	0 minutes at room temperature away from light.			
3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing,			3. Add 2 ml RBC lysis	buffer, incubate for 10 minutes away from light after mixing,			
dissolve red blood cells.			dissolve red blood ce	ells.			
4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the			4. Sample tube is set	to 1000 rpm centrifugation for 5 minutes, discard the			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com