

# **Product Data Sheet**

## Anti-CD103 Antibody-FITC labled

Catalog #	Source	Reactivity	Applications			
CFE8842	Armenian	М	IF, FC			
	Hamster					
Description	Armen	ian Hamster monocl	onal antibody FITC labled to CD103			
Immunogen	Mouse	Mouse intestinal intraepithelial lymphocytes				
Purification	The an	The antibody was purified by affinity chromatography.				
Specificity	Recogr	Recognizes mouse CD103				
Clonality	Monoc	Monoclonal (clone: 2E7)				
Conjugation	FITC	FITC				
Form	Armen	Armenian Hamster IgG. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Dilution	10 µl /	10 μl / assay				
Gene Symbol	ITGAE	ITGAE				
Alternative Na	ames Integri	Integrin alpha-E; HML-1 antigen; Integrin alpha-IEL; Mucosal lymphocyte 1 antigen;				
	CD103					
Entrez Gene	16407	16407 (Mouse)				
SwissProt	Q6067	Q60677 (Mouse)				
<b>Directions for</b>	Use 1. Take	1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
	ml tub	e.				
	2. Add	10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole			
	blood,	incubate for 20 min	utes at room temperature away from light.			
	3. Add	2 ml RBC lysis buffe	; incubate for 10 minutes away from light after mixing,			
	dissolv	e red blood cells.				
	4. Sam	ple tube is set to 10	00 rpm centrifugation for 5 minutes, discard the			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com