

# **Product Data Sheet**

## Anti-CD3e Antibody-PE labled

Catalog #	Source	Reactivity	Applications			
CFF8565	Armenian	Μ	IF, FC			
	Hamster					
Description	Armer	Armenian Hamster monoclonal antibody PE labled to CD3e				
Immunogen						
Purification	The ar	The antibody was purified by affinity chromatography.				
Specificity	Recog	Recognizes mouse CD3e				
Clonality	Mono	Monoclonal (clone: 145-2C11)				
Conjugation	PE	PE				
Form	Armer	Armenian Hamster IgG. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.				
Dilution	10 µl ,	10 μl / assay				
Gene Symbol	CD3E	CD3E				
Alternative N	ames T3E; T	T3E; T-cell surface glycoprotein CD3 epsilon chain; T-cell surface antigen T3/Leu-4				
	epsilo	n chain; CD3e				
Entrez Gene	12501	12501 (Mouse)				
SwissProt	P2264	P22646 (Mouse)				
Directions for	<b>Use</b> 1. Take	1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
	ml tub	ml tube.				
	2. Add	l 10 μl labeled antibo	dy to the bottom of flow tube mixing with the whole			
	blood,	, incubate for 20 min	utes at room temperature away from light.			
	3. Add	2 ml RBC lysis buffe	, incubate for 10 minutes away from light after mixing,			
	dissolv	ve red blood cells.				
	4. San	nple tube is set to 10	00 rpm centrifugation for 5 minutes, discard the			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com