

# **Product Data Sheet**

## Anti-CD96 Antibody-PE labled

Catalog #	Source	Reactivity	Applications		
CFF8616	Mouse	н	IF, FC		
Description		Mouse monoclonal antibod			
Immunogen	-				
Purification		The antibody was purified b	y affinity chromatography.		
Specificity	F	Recognizes human CD96			
Clonality	Ν	Monoclonal (clone: NK92.39			
Conjugation	P	PE			
Form	Ν	Mouse IgG1 kappa. Liquid ir	PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution	1	LO μl / assay			
Gene Symbol	C	CD96			
Alternative Names		T-cell surface protein tactile; Cell surface antigen CD96; T cell-activated increased			
	la	ate expression protein; CD9	6		
Entrez Gene	1	10225 (Human)			
SwissProt	P	240200 (Human)			
<b>Directions for Use</b>		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5			
	n	nl tube.			
	2	2. Add 10 μl labeled antiboo	y to the bottom of flow tube mixing with the whole		
	b	blood, incubate for 20 minu	tes at room temperature away from light.		
	3	3. Add 2 ml RBC lysis buffer,	incubate for 10 minutes away from light after mixing,		
	Ċ	lissolve red blood cells.			
	4	I. Sample tube is set to 100	) rpm centrifugation for 5 minutes, discard the		
	S	supernatant.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com