

Anti-CD61 Antibody-PerCP labeled

| Catalog # | Source | Reactivity | Applications |
|--|--|------------|--------------|
| CFJ8545 | Mouse | H | IF, FC |
| Description | Mouse monoclonal antibody PerCP labled to CD61 | | |
| Immunogen | Native purified human CD61. | | |
| Purification | The antibody was purified by affinity chromatography. | | |
| Specificity | Recognizes human CD61 | | |
| Clonality | Monoclonal (clone: VI-PL2) | | |
| Conjugation | PerCP | | |
| Form | Mouse IgG1. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide. | | |
| Dilution | 10 µl / assay | | |
| Gene Symbol | ITGB3 | | |
| Alternative Names | GP3A; Integrin beta-3; Platelet membrane glycoprotein IIIa; GPIIIa; CD antigen CD61 | | |
| Entrez Gene | 3690 (Human) | | |
| SwissProt | P05106 (Human) | | |
| Directions for Use | <div>1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube.</div> <div>2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light.</div> <div>3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells.</div> <div>4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the supernatant.</div> <div>5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation</div> | | |
| Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference | | | |
| Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish | | | |

COHESION BIOSCIENCES LIMITED

WEB
www.cohesionbio.com

ORDER
order@cohesionbio.com

SUPPORT
techsupport@cohesionbio.com

CUSTOM
custom@cohesionbio.com

Product Data Sheet

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB

www.cohesionbio.com

ORDER

order@cohesionbio.com

SUPPORT

techsupport@cohesionbio.com

CUSTOM

custom@cohesionbio.com