

## **Product Data Sheet**

# Anti-CD36 Antibody-PerCP/Cy5.5 labled

Catalog # Source Reactivity Applications

CFK8584 Mouse H IF, FC

**Description** Mouse monoclonal antibody PerCP/Cy5.5 labled to CD36

Immunogen Native purified human CD36.

**Purification** The antibody was purified by affinity chromatography.

**Specificity** Recognizes human CD36

Clonality Monoclonal (clone: 5-271)

Conjugation PerCP/Cy5.5

Form Mouse IgG2a kappa. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.

Dilution 10 μl / assay

Gene Symbol CD36

Alternative Names GP3B; GP4; Platelet glycoprotein 4; Fatty acid translocase; FAT; Glycoprotein IIIb;

GPIIIB; Leukocyte differentiation antigen CD36; PAS IV; PAS-4; Platelet collagen

receptor; Platelet glycoprotein IV; GPIV; Thrombospondin receptor; CD36

Entrez Gene 948 (Human)

SwissProt P16671 (Human)

Directions for Use 1. Take 100 μl peripheral blood anticoagulated by EDTA and add to the bottom of 5

ml tube.

2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole

blood, incubate for 20 minutes at room temperature away from light.

3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing,

dissolve red blood cells.

4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com





### **Product Data Sheet**

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**