

# **Product Data Sheet**

# Anti-CD200 Antibody-PerCP/Cy5.5 labled

Catalog #	Source	Reactivity	Applications			
CFK8686	Mouse	Н	IF, FC			
Description	I	Mouse monoclonal antiboc	ly PerCP/Cy5.5 labled to CD200			
Immunogen						
Purification	-	The antibody was purified b	by affinity chromatography.			
Specificity	I	Recognizes human CD200				
Clonality	I	Monoclonal (clone: OX-104	)			
Conjugation	I	PerCP/Cy5.5				
Form	I	Mouse IgG1 kappa. Liquid i	n PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.			
Dilution	:	10 μl / assay				
Gene Symbol	(	CD200				
Alternative Names		MOX1; MOX2; OX-2 membrane glycoprotein; CD200				
Entrez Gene		4345 (Human)				
SwissProt	Í	P41217 (Human)				
Directions for Use		1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5				
	I	ml tube.				
		2. Add 10 µl labeled antibo	dy to the bottom of flow tube mixing with the whole			
	I	blood, incubate for 20 minu	utes at room temperature away from light.			
	3	3. Add 2 ml RBC lysis buffer	; incubate for 10 minutes away from light after mixing,			
	(	dissolve red blood cells.				
	4	4. Sample tube is set to 100	00 rpm centrifugation for 5 minutes, discard the			
	9	supernatant.				
	ļ	5. Add 2 ml PBS wash buffe	r to resuspend the cells, then 1000 rpm centrifugation			
A surface to a first			histochemister, IC, lever of histochemist, IC, Class entertain, IC			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com