

# Product Data Sheet

## Anti-CD57 Antibody-AF647 labeled

Catalog #	Source	Reactivity	Applications
CFN8543	Mouse	H	IF, FC
Description	Mouse monoclonal antibody AF647 labled to CD57		
Immunogen	Native purified human CD57.		
Purification	The antibody was purified by affinity chromatography.		
Specificity	Recognizes human CD57		
Clonality	Monoclonal (clone: NK-1)		
Conjugation	AF647		
Form	Mouse IgM. Liquid in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.		
Dilution	10 µl / assay		
Gene Symbol	B3GAT1		
Alternative Names	GLCATP; Galactosylgalactosylxylosylprotein 3-beta-glucuronosyltransferase 1; Beta-1.3-glucuronyltransferase 1; Glucuronosyltransferase P; GlcAT-P; UDP-GlcUA:glycoprotein beta-1.3-glucuronyltransferase; GlcUAT-P		
Entrez Gene	27087 (Human)		
SwissProt	Q9P2W7 (Human)		
Directions for Use	<div>1. Take 100 µl peripheral blood anticoagulated by EDTA and add to the bottom of 5 ml tube.</div> <div>2. Add 10 µl labeled antibody to the bottom of flow tube mixing with the whole blood, incubate for 20 minutes at room temperature away from light.</div> <div>3. Add 2 ml RBC lysis buffer, incubate for 10 minutes away from light after mixing, dissolve red blood cells.</div> <div>4. Sample tube is set to 1000 rpm centrifugation for 5 minutes, discard the</div>		
<b>Application key:</b> E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference			
<b>Species reactivity key:</b> H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish			

### COHESION BIOSCIENCES LIMITED

**WEB**  
www.cohesionbio.com

**ORDER**  
order@cohesionbio.com

**SUPPORT**  
techsupport@cohesionbio.com

**CUSTOM**  
custom@cohesionbio.com

## Product Data Sheet

supernatant.

5. Add 2 ml PBS wash buffer to resuspend the cells, then 1000 rpm centrifugation for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

### Storage/Stability

Shipped and store at 4°C for one year. Do not freeze.

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

#### WEB

[www.cohesionbio.com](http://www.cohesionbio.com)

#### ORDER

[order@cohesionbio.com](mailto:order@cohesionbio.com)

#### SUPPORT

[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

#### CUSTOM

[custom@cohesionbio.com](mailto:custom@cohesionbio.com)