

## **Product Data Sheet**

# Anti-CD56 Antibody-Pacific Blue labled

| Catalog #          | Source | e Reactivity   | Applications   |  |  |
|--------------------|--------|--|--|--|--|
| CFP8542            | Mouse  | e H  | IF, FC   |  |  |
| Description        |        | Mouse monoclonal antibo  | dy Pacific Blue labled to CD56                             |  |  |
| Immunogen          |        | Native purified human CDS  | 56.  |  |  |
| Purification       |        | The antibody was purified  | by affinity chromatography.                                |  |  |
| Specificity        |        | Recognizes human CD56  |  |  |  |
| Clonality          |        | Monoclonal (clone: MY31)   |  |  |  |
| Conjugation        |        | Pacific Blue   |  |  |  |
| Form               |        | Mouse IgG1 kappa. Liquid   | in PBS, pH 7.3, 0.2% BSA, and 0.02% sodium azide.          |  |  |
| Dilution           |        | 10 μl / assay  |  |  |  |
| Gene Symbol        |        | NCAM1  |  |  |  |
| Alternative Na     | ames   | NCAM; Neural cell adhesic  | n molecule 1; N-CAM-1; NCAM-1; CD antigen CD56             |  |  |
| Entrez Gene        |        | 4684 (Human)   |  |  |  |
| SwissProt          |        | P13591 (Human)   |  |  |  |
| Directions for Use |        | 1. Take 100 $\mu l$ peripheral blood anticoagulated by EDTA and add to the bottom of 5 |  |  |  |
|                    |        | ml tube.   |  |  |  |
|                    |        | 2. Add 10 $\mu$ l labeled antibo   | ody to the bottom of flow tube mixing with the whole       |  |  |
|                    |        | blood, incubate for 20 min   | utes at room temperature away from light.                  |  |  |
|                    |        | 3. Add 2 ml RBC lysis buffe  | r, incubate for 10 minutes away from light after mixing,   |  |  |
|                    |        | dissolve red blood cells.  |  |  |  |
|                    |        | 4. Sample tube is set to 10  | 00 rpm centrifugation for 5 minutes, discard the           |  |  |
|                    |        | supernatant.   |  |  |  |
|                    |        | 5. Add 2 ml PBS wash buff  | er to resuspend the cells, then 1000 rpm centrifugation    |  |  |
| Application kow    |        | W/D Western blot III Immun   | histochomistry IE Immunofluorosconco EC Elouy outomotry IC |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



### **Product Data Sheet**

for 5 minutes, discard the supernatant.

6. Add 0.5 ml PBS wash buffer to resuspend the cells and detect by flow cytometry (sample should be determined on the day on the machine and can also be added fixation overnight at 4 °C then measured).

**Storage/Stability** Shipped and store at 4°C for one year. Do not freeze.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com