

Product Data Sheet

Anti-PKC theta Antibody


Catalog #	Source	Reactivity	Applications		
CPA4787	Rabbit	н	WB, IH		
Description		Rabbit polyclonal antibody	to PKC theta		
Immunogen		KLH-conjugated synthetic p	eptide encompassing a sequence within the C-term		
		region of human PKC theta	The exact sequence is proprietary.		
Purification		The antibody was purified b	oy immunogen affinity chromatography.		
Specificity		Recognizes endogenous lev	els of PKC theta protein.		
Clonality		Polyclonal			
Conjugation					
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,			
		and 0.01% sodium azide.			
Dilution		WB (1/500 - 1/1000), IH (1/1	00 - 1/200)		
Gene Symbol		PRKCQ			
Alternative Na	ames	PRKCT; Protein kinase C the	ta type; nPKC-theta		
Entrez Gene		5588 (Human)			
SwissProt		Q04759 (Human)			
Storage/Stabi	lity	Shipped at 4°C. Upon delive	ry aliquot and store at -20°C for one year. Avoid		
		freeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com


Coherion



For research purposes only, not for human use

Product Data Sheet

Western blot analysis of PKC theta expression in MCF7 (A), Jurkat (B) whole cell lysates. (Predicted band size: 81 kD; Observed band size: 78 kD)



Immunohistochemical analysis of PKC theta staining in human skeletal muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com