

Product Data Sheet

MYCBPAP siRNA (Human)

Catalog #	Source	Reactivity		Applications		
CRJ3286	Synthetic	Н		RNAi		
Description	siRN	A to inhibit MYCBPAP e	expression using	RNA interference		
Specificity	MYC	BPAP siRNA (Human) i	s a target-specifi	ic 19-23 nt siRNA ol	igo duplexes designed	
	to kr	nock down gene expres	ssion.			
Form	Lyop	hilized powder				
Gene Symbol	MYC	МҮСВРАР				
Alternative N	ames AMA	AMAP1; MYCBP-associated protein; AMAM-1; AMY-1-binding protein 1; AMAP-1				
Entrez Gene	8407	73 (Human)				
SwissProt	Q8TI	Q8TBZ2 (Human)				
Purity	> 979	> 97%				
Quality Contr	ol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appr	opriate coupling efficie	ency. The oligo is	s subsequently puri	fied by affinity-solid	
	phas	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spec	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the p	previous lot by mass sp	ectrometry to e	nsure maximum lot	-to-lot consistency.	
Components	We c	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	hum	human MYCBPAP gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can b	can be transfected individually or pooled together to achieve knockdown of the				
	targe	target gene, which is most commonly assessed by qPCR or western blot.				
	Cor	nponent	:	15 nmol	30 nmol	
	MY	CBPAP siRNA (Human)	- A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

MYCBPAP siRNA (Human) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

MYCBPAP siRNA (Human) - C	5 nmol x 1	5 nmol x 2	
Negative Control	2.5 nmol x 1	2.5 nmol x 2	
DEPC Water	1 ml x 1	1 ml x 2	

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
96-well		100 nM	0.5 μl	0.25 μl
	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
24-well		100 nM	2.5 μl	1 µl
	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com