

Product Data Sheet

5 nmol x 2

C19orf44 siRNA (Human)

Catalog #	Source	Reactivity	Арр	lications	
CRJ3332	Synthetic	н	RNA	i	
Description	siRNA	to inhibit C19orf44	expression using RNA	interference	
Specificity	C190	C19orf44 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expre	ssion.		
Form	Lyoph	nilized powder			
Gene Symbol	C190	rf44			
Alternative N	lames Unch	aracterized protein C	19orf44		
Entrez Gene	8416	7 (Human)			
SwissProt	Q9H6	5X5 (Human)			
Purity	> 97%	> 97%			
Quality Contr	rol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	opriate coupling effici	ency. The oligo is subs	equently purific	ed by affinity-solid
	phase	e extraction. The anno	ealed RNA duplex is fu	rther analyzed	by mass
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass s	pectrometry to ensure	e maximum lot-t	to-lot consistency.
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human C19orf44 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes			
	can b	can be transfected individually or pooled together to achieve knockdown of the			
	targe	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nm	nol	30 nmol
	C190	orf44 siRNA (Human)	- A 5 nmc	ol x 1	5 nmol x 2
		6			

C19orf44 siRNA (Human) - B 5 nmol x 1 Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

C19orf44 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com