

Product Data Sheet

NUDT16L1 siRNA (Human)

Catalog #	Source	Reactivity		Applications		
CRJ3431	Synthetic	н		RNAi		
Description	siRNA	to inhibit NUDT16L1	expression us	ing RNA interference		
Specificity	NUDT	NUDT16L1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes				
	desig	ned to knock down ge	ene expression			
Form	Lyoph	Lyophilized powder				
Gene Symbol	NUDT	NUDT16L1				
Alternative N	ames SDOS	SDOS; Protein syndesmos; NUDT16-like protein 1				
Entrez Gene	84309	84309 (Human)				
SwissProt	Q9BR	Q9BRJ7 (Human)				
Purity	> 97%	> 97%				
Quality Control Oligonucleotide synthesis is monitored base			se by base through trityl analysis to ensure			
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid				
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the p	revious lot by mass sp	ectrometry to	ensure maximum lot	-to-lot consistency.	
Components We offers pre-designed sets of 3 different target-specific siRNA oligo duplex			A oligo duplexes of			
	huma	human NUDT16L1 gene. Each vial contains 5 nmol of lyophilized siRNA. The				
	duple	duplexes can be transfected individually or pooled together to achieve knockdown				
	of the target gene, which is most commonly assessed by qPCR or western blot.			or western blot.		
	Com	ponent		15 nmol	30 nmol	
	NUD)T16L1 siRNA (Human) - A	5 nmol x 1	5 nmol x 2	
	NUD	T16L1 siRNA (Human) - B	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

N	egative Control	2.5 nmol x 1	2.5 nmol x 2
D	EPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 μl
6-well	2 ml	50 nM	5 μl	5 μl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com