

# **Product Data Sheet**

## SLC25A21 siRNA (Human)

Catalog #	Source	Reactivity	Applications			
CRJ3876	Synthetic	Н	RNAi			
	•					
Description	siRNA	A to inhibit SLC25A21	expression using RNA interfere	ence		
Specificity	SLC25	5A21 siRNA (Human) i	s a target-specific 19-23 nt siRl	NA oligo duplexes designed		
	to kn	ock down gene expre	ssion.			
Form	Lyopł	nilized powder				
Gene Symbol	SLC25	SLC25A21				
Alternative N	ames ODC;	ODC; Mitochondrial 2-oxodicarboxylate carrier; Solute carrier family 25 member 21				
Entrez Gene	8987	4 (Human)				
SwissProt	Q9BC	QT8 (Human)				
Purity	> 97%	> 97%				
Quality Contr	ol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	opriate coupling efficient	ency. The oligo is subsequently	purified by affinity-solid		
	phase	e extraction. The anne	ealed RNA duplex is further and	alyzed by mass		
	spect	rometry to verify the	exact composition of the duple	ex. Each lot is compared to		
	the p	revious lot by mass sp	pectrometry to ensure maximu	m lot-to-lot consistency.		
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	human SLC25A21 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can b	can be transfected individually or pooled together to achieve knockdown of the				
	targe	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	SLC2	25A21 siRNA (Human)	- A 5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

SLC25A21 siRNA (Human) - B

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

DEPC Water 1 ml x 1 1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 μl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
6-well	2 ml	100 nM	10 µl	5 μl
		50 nM	5 μl	5 μl
		10 nM	1 µl	5 μl

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com