

Product Data Sheet

DEFB123 siRNA (Human)

Catalog #	Source	Reactivity		Applications		
CRJ6608	Synthetic	Н		RNAi		
Description	siRNA	to inhibit DEFB123 e	xpression using	g RNA interference		
Specificity	DEFB1	DEFB123 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed				
	to kno	ock down gene expres	sion.			
Form	Lyoph	ilized powder				
Gene Symbol	DEFB1	DEFB123				
Alternative N	ames DEFB2	DEFB23; Beta-defensin 123; Beta-defensin 23; DEFB-23; Defensin beta 123				
Entrez Gene	24593	245936 (Human)				
SwissProt	Q8N68	Q8N688 (Human)				
Purity	> 97%	> 97%				
Quality Contr	ol Oligon	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	priate coupling efficie	ency. The oligo	is subsequently pur	ified by affinity-solid	
	phase	extraction. The anne	aled RNA duple	ex is further analyze	ed by mass	
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	evious lot by mass sp	ectrometry to	ensure maximum lo	ot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	humai	n DEFB123 gene. Each	n vial contains !	5 nmol of lyophilize	d siRNA. The duplexes	
	can be	e transfected individu	ally or pooled t	together to achieve	knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent		15 nmol	30 nmol	
	DEFB	123 siRNA (Human) -	A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

DEFB123 siRNA (Human) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Ν	legative Control	2.5 nmol x 1	2.5 nmol x 2
C	DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com