

Product Data Sheet

SLC26A5 siRNA (Human)

Catalog #	Source	Reactivity		Applications		
CRJ7676	Synthetic	н		RNAi		
Description	siRN	A to inhibit SLC26A5 e	xpression using	RNA interferen	ce	
Specificity	SLC2	SLC26A5 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed				
	to kr	nock down gene expres	ssion.			
Form	Lyop	hilized powder				
Gene Symbol	SLC2	SLC26A5				
Alternative N	lames PRES	PRES; Prestin; Solute carrier family 26 member 5				
Entrez Gene	3756	511 (Human)				
SwissProt	P587	743 (Human)				
Purity	> 979	> 97%				
Quality Cont	rol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appr	opriate coupling efficient	ency. The oligo i	is subsequently	purified by affinity-solid	
	phas	phase extraction. The annealed RNA duplex is further analyzed by mass				
	spec	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the p	previous lot by mass sp	pectrometry to	ensure maximur	n lot-to-lot consistency.	
Components	We c	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	hum	human SLC26A5 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can b	can be transfected individually or pooled together to achieve knockdown of the				
	targe	target gene, which is most commonly assessed by qPCR or western blot.				
	Con	nponent		15 nmol	30 nmol	
	SLC	26A5 siRNA (Human) -	- A	5 nmol x 1	5 nmol x 2	

SLC26A5 siRNA (Human) - B5 nmol x 15 nmol x 2Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-

Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

SLC26A5 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 μl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 μl	5 µl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com