

Product Data Sheet

Recombinant Human Noggin Protein

Catalog # Source Reactivity Applications

CRP2780 Human cells Human E, WB, SDS-PAGE, MS

Description Recombinant Human Noggin Protein is produced by mammalian expression system

and the target gene encoding Gln28-Cys232.

Form Lyophilized from a 0.2 μm filtered solution of 20mM PB, 500mM NaCl, 2mM EDTA,

pH7.4.

Gene Symbol NOG

Alternative Names Noggin

Entrez Gene 9241 (Human)

SwissProt Q13253 (Human)

Purity Greater than 95% as determined by reducing SDS-PAGE.

Chemical Structure QHYL HIRPAPSDNL PLVDLIEHPD PIFDPKEKDL NETLLRSLLG GHYDPGFMAT

SPPEDRPGGG GGAAGGAEDL AELDQLLRQR PSGAMPSEIK GLEFSEGLAQ GKKQRLSKKL

RRKLQMWLWS QTFCPVLYAW NDLGSRFWPR YVKVGSCFSK RSCSVPEGMV CKPSKSVHLT

VLRWRCQRRG GQRCGWIPIQ YPIISECKCS C

Quality Control Endotoxin: Less than 0.1 ng/μg (1 IEU/μg) as determined by LAL test.

Directions for Use Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not

recommended to reconstitute to a concentration less than 100 µg/ml. Dissolve the

lyophilized protein in ddH2O. Please aliquot the reconstituted solution to minimize

freeze-thaw cycles.

Storage/Stability Lyophilized protein should be stored at -20°C, though stable at room temperature

for 3 weeks. Reconstituted protein solution can be stored at 2-8°C for 2-7 days.

Aliquots of reconstituted samples are stable at -20°C for 3 months.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com