

Product Data Sheet

Goat Anti-Human IgG gamma chain - TRITC

Catalog #	Sourc	e Reactivity	Applications	
CSA2605	Goat	Н	E, IH, IF, FC	
Description		Goat Polyclonal Secondary Antibody to Human IgG gamma chain TRITC labeled		
Immunogen		Human IgG gamma chain		
Purification		The antibody was isolated from antisera by immunoaffinity chromatography using		
		antigens coupled to agaro	se beads.	
Specificity		By immunoelectrophoresis and ELISA this antibody reacts specifically with Human		
		IgG gamma chain. No anti	body was detected against non immunoglobulin serum	
		proteins.		
Clonality		Polyclonal		
Conjugation		TRITC		
Form		1 mg/ml. Liquid in 0.01M	Phosphate Buffered Saline, pH 7.2, containing 1% BSA,	
		50% glycerol, 0.02% Sodiu	m Azide	
Dilution		IH (1/100 - 1/1000), IF (1/10	00 - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay	
		dependent concentration		
Fluorophore		Ex/Em = 595/613 nm		
Storage/Stabi	ility	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid		
		freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com