

Product Data Sheet

Goat Anti-Rabbit IgG (H&L) - TRITC

Catalog #	Source	e Reactivity	Applications		
CSA2611	Goat	Rb	E, IH, IF, FC		
Description		Goat Polyclonal Secondary Antibody to Rabbit IgG (H&L) TRITC labeled			
Immunogen		Rabbit IgG			
Purification		The antibody was isolated from antisera by immunoaffinity chromatography using			
		antigens coupled to agaros	e beads.		
Specificity		By immunoelectrophoresis and ELISA this antibody reacts specifically with Rabbit			
		IgG. No antibody was deter	cted against non immunoglobulin serum proteins.		
Clonality		Polyclonal			
Conjugation		TRITC			
Form		1 mg/ml. Liquid in 0.01M P	hosphate Buffered Saline, pH 7.2, containing 1% BSA,		
		50% glycerol, 0.02% Sodiur	n Azide		
Dilution		IH (1/100 - 1/1000), IF (1/100) - 1/1000), FC (1/1000 - 1/4000), E (Use at an assay		
		dependent concentration)			
Fluorophore		Ex/Em = 595/613 nm			
Storage/Stabi	lity	Shipped at 4°C. Upon delive	ery aliquot and store at -20°C for one year. Avoid		
		freeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com